

For submission to: *Bioengineering and Translational Medicine*

Early nanoparticle intervention preserves motor function following cervical spinal cord injury

Sarah E. Hocevar, Brian C. Ross, Yinghao Wang, Cecelia R. Crowther, Samantha R. Schwartz, Brain J. Cummings, Aileen J. Anderson, Lonnie D. Shea*

Affiliations:

Neuroscience Graduate Program, University of Michigan Medical School, Ann Arbor, MI, USA
204 Washtenaw Ave. Ann Arbor, MI 48109, USA

S.E. Hocevar, L.D. Shea

Department of Biomedical Engineering, University of Michigan, USA
2200 Bonisteel Blvd, Ann Arbor, MI 48109, USA

S.E. Hocevar, B.C. Ross, Y Wang, C.R. Crowther, S.R. Schwartz, L.D. Shea

Department of Chemical Engineering, University of Michigan, Ann Arbor, MI, USA
2300 Hayward St, Ann Arbor, MI 48109, USA

L.D. Shea

Institute for Memory Impairments and Neurological Disorders, University of California, Irvine, California

Biological Sciences III, 2642, Irvine, CA 92697

A.J. Anderson, B.J. Cummings

Sue and Bill Gross Stem Cell Research Center, University of California, Irvine, California
845 Health Sciences Rd, Irvine CA 92697

A.J. Anderson, B.J. Cummings

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](https://doi.org/10.1002/btm2.70011). Please cite this article as doi: [10.1002/btm2.70011](https://doi.org/10.1002/btm2.70011)

Physical Medicine and Rehabilitation, University of California, Irvine, California

18124 Culver Dr # F, Irvine, CA 92612

A.J. Anderson, B.J. Cummings

*Corresponding author

ldshea@umich.edu

(734) 647-6319

3318 NCRC Building 520

1600 Huron Parkway

Ann Arbor, MI 48109, USA

Abstract

Spinal cord injury (SCI) triggers an immediate influx of immune cells that secrete pro-inflammatory cytokines and reactive oxygen species that cause tissue damage that is secondary to the initial physical trauma. We aim to reprogram these immune cells to promote a less inflammatory and more pro-regenerative environment. Herein, we investigated the window in time which poly(lactide-co-glycolide) nanoparticles (NPs) administration can successfully modulate the immune response and promote functional sparing. The dynamics of immune cell infiltration and secondary tissue damage were studied following injection of NPs intravenously every 24 hours for 7 days following injury with the first injection starting at 2-, 4-, or 24-hours post-injury (hpi). At 7 days post-injury (dpi), early NP intervention decreased the number of infiltrating macrophages and neutrophils but delaying treatment until 24 hpi increased the number of neutrophils above control. All mice that received NPs had greater neuronal sparing contralateral to the injury, but mice that received NPs at early timepoints have greater neuromuscular junction innervation and motor endplate sparing. The increased sparing of neurons and neural circuits in the 2 hpi NP group corresponded with increased motor function, as measured by a ladder beam test. Collectively, these results suggest that early intervention with NPs can modulate the inflammatory response and preserve motor function and circuits following SCI.

Translational Impact Statement

Inflammation following spinal cord injury causes secondary damage to neural tissue and exacerbates function loss. There are currently no standard-of-care treatments that address inflammation after spinal cord injury. This study describes the use of drug-free, biodegradable poly(lactide-co-glycolide) nanoparticles to acutely reprogram peripheral immune cells following spinal cord injury (SCI), which can reduce inflammation and the associated damage and lead to improved functional recovery.

Keywords: spinal cord injury; immunomodulation; nanoparticle; inflammatory response

Introduction

The United States alone has approximately 17,000 new cases of spinal cord injury (SCI) each year ¹. Following injury, circulating neutrophils infiltrate the spinal cord within hours of injury and release inflammatory cytokines and reactive oxygen and nitrogen species at the injury that exacerbate neuronal and oligodendrocyte cell death ²⁻⁴. Methylprednisolone, a glucocorticoid that downregulates adhesion molecule and cytokine production in immune cells, has been investigated in clinical trials to decrease the inflammatory response to SCI ⁵⁻⁷. However, while some patients exhibited improved motor and sensation function, its use has been discontinued because of its many off-target effects including sepsis, pneumonia, impaired wound healing, and gastrointestinal bleeding. Additionally, mouse studies have shown that complete ablation of neutrophils and macrophages does not significantly improve motor performance and has a negative effect on wound healing and regrowth into the injury ⁸⁻¹⁰. Together, these studies demonstrate the need for a clinical intervention that can attenuate immune cells during spinal cord injury without causing widespread immunosuppression and its negative effects.

An emerging strategy for SCI to address inflammation is polarizing innate immune cells toward a less-inflammatory phenotype, which can lead to phenotypes that promote rather than diminish functional recovery. Biomaterial technologies in particular have emerged as tools to locally modulate inflammation and enhance the regenerative potential of the spinal cord. Recently nanoparticles (NPs) have been reported to target spinal cord injury due to their ability to cross the semipermeable blood-brain barrier ^{11,12}. Nanoparticles are often conjugated to a therapeutic agent and administered systemically, yet limited accumulation of NPs at the injury limits the extent of the therapeutic effect. Conjugating additional guiding peptides onto the NPs can increase accumulation in the spinal cord, but the therapeutic effect can still be limited by the low phagocytic ability of many resident cells of the spinal cord ¹³⁻¹⁶. To circumvent these accumulation issues and the off-target effects of drugs like methylprednisolone, our lab has previously developed cargo-

less poly(lactide-co-glycolide) (PLG) NPs that are phagocytosed by circulating myeloid cells and reprograms them toward a more anti-inflammatory phenotype. Administration of these NPs led to increased axon regrowth into the injury and improved functional recovery in mice as early as 7 days post injury that were sustained for at least 12 weeks ¹⁷.

In this report, we investigate the cellular processes that are influenced by NP administration, with assessment of the immune dynamics following administration, and sparing of neural tissue. The studies employ a cervical hemisection injury, and the timing of initial NP administration was varied between 2 and 24 hours, with daily NP injections subsequently for one week. This model was chosen instead of the more common contusion model due to more severe functional and behavioral outcomes following penetrating injuries. The varied timing of first NP administration was anticipated to target different populations of infiltrating immune cells. An administration time of 2 hours would target neutrophil and monocyte populations that are upregulated following injury, whereas a 4-hour time point would preferentially target monocytes ^{18,19}. An administration time of 24 hours was chosen because both neutrophils and monocytes have already begun to accumulate in the injury ²⁰. Because this study is a continuation of our previous work, we elected to use the same injury model which includes the implantation of a biomaterial bridge that limits scar formation and supports axon growth into the injury. The presence of the bridge, however, increases the permeability of the injury and NPs can accumulate in the injury without being transported by immune cells.²¹ While the numbers of infiltrating immune cells are similar in NP-treated and control mice 2 days post-injury, fewer cells were present by the end of the first week post-injury. All NP conditions led to increased survival of neurons in the spinal cord compared to controls, but earlier intervention improved neuromuscular junction innervation (NMJ) and motor performance. Collectively, these results support the use of NPs as a means to modulate the acute inflammatory response to improve outcomes in spinal cord injury.

Methods

PLG NP fabrication and administration

PLG NPs were made by dissolving 50:50 PLG (inherent viscosity = 0.55dL/g; Evonik, Sand Creek, MI) in dichloromethane along with poly(ethylene-alt-maleic anhydride) (Polysciences, Inc., Warrington, PA). The mixture was sonicated with a Cole-Parmer CPX130 Ultrasonic Processor (Vernon Hills, IL) and then stirred overnight to evaporate the organic phase. The following day the NPs were collected and washed with water to produce particles that are 626.95 ± 47.67 nm in size (average pdi = 0.19) and -47.86 ± 4.38 mV in charge. Particle size and charge were measured with a Zetasizer Nano ZSP (Malvern Panalytical, Malvern, United Kingdom). The particles lyophilized with sucrose and D-mannitol (Sigma-Aldrich, St. Louis, MO). During administration, each mouse received 0.1 mL of 10 mg/mL NPs in DPBS injected into the lateral tail vein daily for one week following injury. Control mice received 0.1 mL DPBS instead of NPs. The timing of the first NP injection varied from 2 hours post-injury (hpi) up to 24 hpi. Every subsequent NP injection was 24 hours after the previous injection so that NP injections remained staggered between different timing conditions.

Multichannel PLG bridge fabrication

Bridges were fabricated using the same protocol previously published by our lab²²⁻²⁴. PLG (75:25 lactide:glycolide; inherent viscosity = 0.76 dL/g; Lakeshore Biomaterials, Birmingham, AL, USA) was dissolved in dichloromethane (6 % w/w) and emulsified in 1% poly (vinyl-alcohol) (Millipore Sigma, Burlington, MA) using a homogenizer (PolyTron 3100; Kinematica AG, Littau, Switzerland). The resulting microspheres were approximately 1 μ m in diameter. D-sucrose (Millipore Sigma), D-glucose (Millipore Sigma) and dextran MW 100,000 (Millipore Sigma) were mixed at a ratio of 5.3:2.5:1, respectively, by mass and then were caramelized, cooled, and drawn from solution using a Pasteur pipette to create sugar strands. Each 150-250 μ m strand

was coated with a 1:1 mixture of PLG microspheres and salt (63–106 μ m) and pressed into a salt-lined aluminum mold. The strands were used to create channels to act as a conduit for regenerating axon bundles and salt was used to create a microporous structure. The materials were gas-foamed under 800 psi for 16 hours to fuse the particles into a matrix. The bridges were cut into 1.15 mm long pieces and the NaCl was leached with water, leaving behind a multichannel bridge.

Spinal cord hemisection model and animal care:

All animal work was performed according to protocol number PRO00011456 approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Michigan. A C5 lateral hemisection was performed on 6-8 week old male and female C57BL/6J (Jackson Laboratory, Bar Harbor, ME) or Crym-RFP (cross between Mutant Mouse Resource & Research Center #036627-UCD and Jackson Laboratory #007905) mice as previously described²². C57BL/6J mice were only used in flow cytometry experiments; all behavioral and histology experiments used Crym-RFP mice that have been used in our previous studies. Mice were anesthetized with 2% isoflurane and 1 mg/kg bupivacaine was administered as a local anesthetic. After a C5 laminectomy was performed, a 1.15 mm lateral hemisection of the left side of the spinal cord was removed to allow for PLG bridge implantation. The injury and exposed spinal cord were covered in Gelfoam (Ethicon, Raritan, NJ). Muscles were sutured using 5/0 Chromic Gut (Henry Schein, Melville, NY), and the skin was stapled. Post-operative care consisted of buprenorphine (0.1 mg/kg) twice daily for 3 days, enrofloxacin (2.5 mg/kg, Bayer, Leverkusen, Germany) daily for 14 days, and lactated ringers' solution (1 mL, MWI Veterinary Supply, Boise, ID) daily for 5 days. Bladders were expressed twice daily until function recovered and wound clips were removed 10 days after surgery.

Behavioral analysis

The horizontal ladder beam test was used to evaluate locomotor function over a period of 12 weeks following injury. Mice were trained on the ladder beam consisting of 50 rungs during the week prior to injury. Mice were recorded at 1, 2, and every two weeks up to 12 weeks following injury. Videos were scored by two blinded observers for 3 trials per animal by counting the number of correct and misplaced steps of the left forepaw on the ladder beam.

Immunohistochemistry

At 1-week post-injury, mice were euthanized with 5% isoflurane and C4-C6 along with the left acromiotrapezius and forearm flexor muscles were dissected. Tissues were flash frozen in isopentane on dry ice and embedded in 20% w/w sucrose (Millipore Sigma) in OCT (ThermoFisher Scientific). Muscles and spinal cords were cryosectioned at 14 μ m thick longitudinally and transversely, respectively. Tissue sections were permeabilized in 0.5% Triton in Tris-buffered saline, fixed with 4% paraformaldehyde, and blocked with normal goat or normal donkey serum as necessary.

Antibodies used were mouse anti-NeuN (1:250, Millipore, Burlington, MA, #MAB377) for mature neurons, chicken anti-NF200 (1:250, AVES Labs, Davis, CA, #NFH) for neurofilaments, anti-bungarotoxin conjugated to Alexa Fluor 647 (1:500, Thermo Fisher, Waltham, MA, #B35450) for motor endplates, rat anti-F4/80 (1:200, Bio-rad MCA497GA) for macrophages, and rat anti-Ly6G (Biolegend #127602) for neutrophils. Secondary antibodies were AlexaFluor 555 goat anti-rat (1:1000; Invitrogen #A21434), AlexaFluor 555 goat anti-mouse (1:1000; Invitrogen #A21424), and AlexaFluor 488 goat anti-chicken (1:1000; Invitrogen #A11039). Tissues were imaged on an Axio Observer Z1 using an ORCA-Flash 4.0 V2 Digital CMOS camera.

Quantification of IHC

For immune cell and neuron IHC, two slides with four tissue sections were used per mouse.

Tissue sections spanned the length of the C5 injury. Tissue sections above or below the level of injury were not used. For NMJ IHC, two slides with six tissue sections spanning the depth of the muscle were used per mouse. To quantify immune cells in the spinal cord, images were exported as .CZI files and converted to .H5 files in CellProfiler ²⁵. The injury area was outlined using the polygon tool in Fiji and saved as a mask ²⁶. The .H5 files were uploaded to Ilastik and pixel classification was performed after training a set of 4 randomly selected tissues per condition ²⁷. A single operator trained Ilastik to ensure consistent staining identification, and the program was trained against a no primary control to prevent identification of non-specific staining. Immune cells were identified by colocalization of greater than 3 pixels of either F4/80 for macrophages/microglia or Ly6G for neutrophils with DAPI. Only immune cells in the injury area were counted.

The number of NeuN+ neurons on the C5 uninjured side of the spinal cord contralateral to the injury were manually counted by two independent, blinded observers using Fiji. BGT+ motor endplates and innervated motor endplates, defined as colocalization of NFM+ axons with BGT+ motor endplate, were manually counted by two independent, blinded observers using Fiji. The entire longitudinal muscle section was quantified using a total of 12 tissue sections per mouse. The polygon tool in Fiji was used to outline the muscle section and measure the area to determine NMJ density.

Flow cytometry

At 2, 5, or 7 dpi, mice were euthanized with 5% isoflurane and C4-C6 of the spinal cord was dissected, including both the injury site and contralateral side. For uninjured controls, C4-C6 was also dissected. Trypsin (0.5 mg; Millipore Sigma) and collagenase (1 mg; Millipore Sigma) in DMEM were added to a tube with the spinal cords and the spinal cords were minced with microscissors. Following mincing, tissues were incubated in at 37 C in a rotisserie shaker for 20

minutes. The samples were spun down in a microcentrifuge and neutralized with DMEM with 10% FBS (Thermofisher Scientific). The samples were spun down again and supernatant aspirated. Then the tissues were myelin depleted using the Miltenyi Myelin Removal Beads II kit (Bergisch Gladbach, Germany). Following myelin depletion, cells were counted and 1 million cells per animal were used for flow cytometry. Fixable Live/Dead Violet was used to assess cell viability. Samples were incubated with anti-CD16/32 (1:100, clone 93; Biolegend) to block nonspecific staining. Cells were additionally incubated with antibodies for CD11b BV421 (0.2 ug/uL; Biolegend 101251) Ly6C BV510 (0.1 ug/uL; Biolegend 128033), Ly6G PE (0.2 ug/uL; Biolegend 127608), F4/80 AF647 (0.5 ug/uL; Biolegend 157314), and CD45 FITC (0.5 ug/uL; Biolegend 103108) and analyzed using a BioRad ZE5 Cell Analyzer. Data analysis was performed using FlowJo (BD).

Statistical Analysis

Statistical analysis was performed in Graphpad Prism 10. All values are expressed as the mean $\pm$ standard error of the mean. For flow cytometry n=5-6 mice were used. For immunohistological data, n=4-10 mice were used. Number of mice used in a specific figure is noted in the figure caption. For flow cytometry and immunohistochemistry, an ordinary one-way ANOVA was used with Tukey's post-hoc test to correct for multiple comparisons. For ladder beam data, n=10 mice were used and a two-way ANOVA with Tukey's post-hoc test was performed. The p-values for statistical significance are represented with stars (Unless otherwise noted: *p < 0.05, **p < 0.005, ***p < 0.0005, ****p < 0.0001). While both male and female mice were used, studies were not sufficiently powered to analyze sex differences.

Results

Immune cell dynamics across first week of injury

We first analyzed the immune cell response in the spinal cord throughout the first week of injury with and without NP intervention. Our previous study using NPs in a hemisection model of SCI showed differences in immune cell accumulation at the injury at 1 week post-injury; therefore, we wanted to further explore the inflammation dynamics at earlier timepoints in the first week of injury.¹⁷ Administered NPs were 626.95 ± 47.67 nm in size (average pdi = 0.19) and -47.86 ± 4.38 mV in charge, which were similar in size and charge to NPs used in previous studies by our lab.^{17,28} Mice received NPs or PBS starting within 2 hours post-injury (hpi) and received daily NP injections. Flow cytometry for immune cells was performed at 2, 5, or 7 days post-injury (dpi). NP administration trended toward a lower percentage of total CD45+ cells in the spinal cord compared to the PBS control at 5 dpi ($p = 0.0701$) (Figure 1A). The percentage of CD11b+ myeloid cells significantly increased across the first week for PBS-treated mice ($p = 0.0116$), yet not for NP-treated mice (Figure 1B). At 5 dpi, the percentage of myeloid cells was significantly lower in mice that received NPs ($p = 0.0222$). Within each condition, the percentage of F4/80+ cells (macrophages and microglia) increased across the week, and by day 7, significantly fewer F4/80+ cells were observed in mice that received NPs ($p = 0.0207$) (Figure 1C). The percentages of Ly6G+ neutrophils (Figure 1D) and Ly6C+ monocytes (Figure 1E) decrease across the first week within each condition and at day 5 are lower in NP-treated mice. In total, these results show that NPs are able to decrease the accumulation of innate immune cells during the first week after injury.




Figure 1. Immune cell population dynamics in spinal cord across first week of injury.

Mice were given daily injections of NPs or PBS starting at 2 hours post-injury. Spinal cords were harvested at multiple timepoints across the first week of injury. Spinal cords were disassociated and myelin depleted before flow cytometry for immune cell surface markers for total immune cells (A), myeloid cells (B), macrophages and microglia (C), neutrophils (D), and monocytes (E). Data are mean $\pm$ SEM. $n=5-6$ mice per condition. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$

Nanoparticle administration alters immune infiltration into the spinal cord

We investigated the efficacy of NPs when the initial administration occurs at distinct time points following injury, which may reflect the delays in treatment that may be present in a clinical situation. NP were initially administered at 2-, 4-, and 24 hpi, with daily administration until one week. Timing of nanoparticle administration remained staggered throughout the entire week to retain administration timing differences between groups. At 7 dpi, spinal cords were harvested, and immune cell infiltration was quantified by immunohistochemistry for F4/80+ macrophages/microglia (Figure 2A) and Ly6G+ neutrophils (Figure 2B). Consistent with our previous studies, NP administration starting within 2 hpi trended toward a decrease in the number of F4/80+ macrophages/microglia and Ly6G+ neutrophils compared to PBS controls. However, these results were not significantly different. While F4/80+ macrophages were significantly lower

at 7 dpi in NP 2 hpi vs PBS mice using flow cytometry in Figure 1C, neutrophil and total immune cell percentages were not significantly lower, which explains the lack of significant difference using IHC quantification. Interestingly, delayed NP injections starting at 4 and 24 hours resulted in higher Ly6G+ neutrophil infiltration than the PBS control (NP 24 hpi vs PBS: $p = 0.0072$) and early NP intervention (NP 24 hpi vs NP 2 hpi: $p=0.0002$, NP 4hpi vs NP 2 hpi: $p = 0.0022$). This result may be the result of NP administration being too late to effect neutrophil migration to the spinal cord, which typically starts within 3 hours of injury.




Figure 2. Immune cell infiltration in spinal cord at 7 dpi. Mice received daily injections of NPs or PBS control starting at varying times within first day of injury through intravenous injection for one week. Spinal cords were harvested at 7 dpi and IHC for macrophages (A) and neutrophils (B) was performed. Data are mean $\pm$ SEM. $n=4-9$ mice per condition. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

NP administration enhances neuron sparing in spinal cord

We next hypothesized that NP delivery has the ability to influence neuron sparing within the injured spinal cord. Following spinal cord injury, inflammation and excitotoxicity resulting from the trauma and inflammatory cytokine release can induce widespread neuron death. Immunohistochemistry was performed at 1 wpi for NeuN+ mature neurons within the contralateral side to the injury, where inflammatory cytokine concentration should be highest (Figure 3A, circled in yellow). All injured conditions had fewer neurons than uninjured controls, and NP administration starting at 2- and 4- hours post-injury trended toward increased neuron count compared to the PBS control ($p = 0.0521$ and 0.0145) (Figure 3B).




Figure 3. Neuron quantification contralateral to injury at 7 dpi. Mice received daily injections of NPs or PBS control starting at varying times within first day of injury through intravenous injection for one week. Spinal cords were harvested at 7 dpi and IHC was performed for neurons (A). Neurons contralateral to the C5 hemisection were quantified (B). Data are mean $\pm$ SEM. $n = 5-7$ mice per condition. * $p < 0.05$, *** $p < 0.001$, **** $p < 0.0001$

NMJ sparing is enhanced by early NP administration

We next investigated the correlation between NP administration and the sparing of neural circuits. These circuits were quantified in the muscle of the acromiotrapezius (ATX) and forearm flexor muscles through IHC for bungarotoxin+ (BGT+) motor endplates and neurofilament 200+ axons (NFM+) (Figure 4A). SCI can cause the death of lower motor neurons, causing the denervation of NMJs. Prolonged loss of synaptic input leads to the disintegration of orphaned motor endplates and muscle atrophy²⁹⁻³². The ATZ is innervated from C5 and therefore directly affected by the injury (Figure 4B-D), whereas the flexor muscles are innervated from C6 below the injury (Figure 4E-G)³³. In the ATZ, NP treatment did not significantly increase sparing of NMJs compared to PBS controls. However, in the flexor muscles, early NP treatment starting at 2 hpi had a significantly greater percentage of innervated motor endplates compared to control ($p=0.0004$). Delayed NP treatment starting at 4 and 24 hpi lead to decreased sparing of motor endplates and innervated NMJs in both the ATZ and flexor muscles. This may be an effect of the increased infiltration of neutrophils and macrophages at 1 wpi in these conditions. Overall, early NP treatment may not be able to spare NMJ innervation that is associated with the hemisection, but it can help spare innervation caudal to the injury site.




Figure 4. Neuromuscular junction innervation at 7 dpi. Mice received daily injections of NPs or PBS control starting at varying times within first day of injury through intravenous injection for one week. Muscles were harvested at 7 dpi and IHC was performed for axons (neurofilament medium chain, NFM) and motor endplates (bungarotoxin, BGT). (A). NMJs were quantified

from the acromiotrapezius muscle (B-D) and flexor carpi ulnaris and flexor digitorum profundus (E-G). Data are mean $\pm$ SEM. n = 5-10 mice per condition. *p < 0.05, **p < 0.01, *** p < 0.001, ****p < 0.0001

Early NP administration increases locomotor functional sparing

Mice underwent a horizontal ladder beam test to investigate whether the spared neurons and NMJs in the NP conditions correlated with better locomotor function. At 7 dpi, mice that received NPs within 4 hpi performed significantly better than PBS controls with more correct placements (Figure 5A) and fewer missed rungs (NP 2 hpi vs PBS: p = 0.0243, NP 4 hpi vs PBS: p = 0.0491) (Figure 5B). Mice that did not receive NPs until 24 hpi performed similarly to mice that only received PBS in number of misses. After 7 dpi, the motor performance of the NP 2 hpi condition remained steady, but the late NP condition improved over 6 weeks. The performance of the PBS group did not significantly change over the course of the experiment. Collectively, NP administration improved locomotor recovery during the acute and subacute phases of injury if administered within 4 hours after SCI.




Figure 5 Early NP treatment improves ladder beam performance. The horizontal ladder beam test was performed weekly following injury. The number of correct placements (A) and incorrect placements (B) were counted. Data are mean $\pm$ SD. n = 10 per condition. *p < 0.05, *** p < 0.001, ****p < 0.0001

Discussion

In these studies, we investigated the use of NPs to reprogram circulating immune cells to enhance regeneration and functional recovery after penetrating spinal cord injury. While contusion and compression injuries are more common, patients with penetrating injuries experience little spontaneous recovery after injury and there are no clinical therapies designed to reconnect severed spinal cord tracts.^{34,35} SCI leads to the activation of innate immune cells, such as inflammatory monocytes and neutrophils, in the blood and spleen, which can subsequently traffic to the injured spinal cord^{36,37}. This infiltration into the SCI can induce damage secondary to the initial injury, thereby contributing to neuronal cell death, axonal dieback, demyelination, and scar tissue formation^{18,38}. Studies that have attempted depletion of these circulating cells have indicated worsened histological and functional outcomes⁸. NPs, in contrast, do not function to deplete immune cells and instead reprogram their function. After injury, the activated innate immune cells, which have high phagocytic capacity for scavenging cell debris and foreign materials, can internalize the NPs while in circulation. This internalization of the NPs can alter the trafficking and functional phenotype of the innate cells²⁸, which has led to improved functional recovery¹⁷. The immune response following SCI can be highly dynamic, and herein we investigated the timing of NP administration.

NP administration, regardless of the timing for the initial administration, had a different immune cell composition in the spinal cord at 1 week post injury. Previous reports had indicated that NPs delivered at 2 hours post injury decreased the number of inflammatory monocytes and neutrophils at 1 week post injury¹⁷. Herein, we observed a decreased number of macrophages within the injury for all mice that received nanoparticles. Interestingly, mice that received NPs within 4 hours of injury also had fewer neutrophils in the spinal cord, yet mice receiving particles starting at 24 hpi had greater accumulation of neutrophils at 1 week relative to control. Analyzing responses over time, only NP administration at 2 hpi resulted in a smaller percentage of myeloid

cells at 2, 5, and 7 dpi. The immune cell dynamics play an important role in the response that leads to their repair or regeneration.

NP injections within 4 hours of injury yielded greater preservation of motor function at 1 wpi relative to later administration, with retention of this increased function through day 42. This is consistent with human clinical trials of the immunosuppressant methylprednisolone. When patients received a 24-hour long infusion of methylprednisolone within 8 hours of injury, mean motor scores improved at 6 and 12 months post-injury.³⁹ However, when methylprednisolone was started later than 8 hours post-injury there was no motor improvement compared to controls and this was performed for 24 hours in a hospital setting, whereas NPs could be administered by EMTs and NP administration takes mere minutes. Inflammatory cytokines and reactive oxygen species released by pro-inflammatory neutrophils and macrophages after SCI are cytotoxic towards neuronal cells^{40,41}. IL-1 α and TNF α in the injury microenvironment induce increased trafficking of glutamate receptors to neuron cell membranes and thereby cause increased excitotoxicity^{42,43}. NP administration is able to alter the immune cell accumulation and polarization normally seen following SCI and contribute to the increased function by increasing sparing of neurons and motor circuits. Previous studies with NP injection had been performed with a thoracic injury, and BMS scores for NP treated were increased relative to control mice by day 7¹⁷. For the study herein that had a cervical injury, an increased functional response was again observed as early as day 7 for mice receiving particles at 2 hpi relative to control. At 1 wpi, all NP-treated mice, regardless of the timing for initial NP administration, had a larger number of surviving neurons contralateral to the injury than mice that did not receive NPs. Mice that received NPs starting at 24 hpi had fewer intact NMJs than even control mice receiving PBS injections at 2 hpi. While we used both male and female mice in equal number in this study, the number of animals used was not sufficiently powered to study sex-based differences in immune response and neural sparing. This may be significant due to male and female mice exhibiting different immune responses following injury, with male mice exhibiting worse microglia accumulation in the injury and worse

neuron-sparing as a result.^{44,45} Our results suggest that even though neuron survival is not dependent upon time of first NP administration, early administration of NPs may maintain neuronal connections more effectively than later administration.

Mice with a spinal cord injury that received NPs at 24 hpi initially had motor performance similar to control mice, yet their motor performance improved over time. While unchecked immune response after SCI can lead to neuron cell death and axonal dieback, an attenuated inflammatory response can have reduced damage, such as demyelination of nerve fibers without axonal death, which leads to decreases in their conductive potential and functional deficits^{46,47}. After the acute phase of injury, the surviving nerve fibers may have conduction recover partially or completely⁴⁷. For the mice receiving particles later, we hypothesize that the initial deficit and gradual improvement may be associated with recovery from the conduction block. These mice had greater neuronal and NMJ sparing than mice that receive NPs at later times; thus, the recovery in function that was observed during subacute and chronic phases may reflect partial recovery in conduction in the spared motor circuits. Within the mice receiving NPs at 24 hpi, we anticipate that the recovery of function was associated with this recovery in the conduction of spared circuits. However, the initial immune response decreased the extent of spared NMJs relative to NPs delivered at earlier times, and thus the extent of recovery was more modest.

In conclusion, this report demonstrates that NP treatment can modulate the inflammatory microenvironment after SCI, and that NP administration within 24 hours is associated with a greater recovery of function relative to control. While the numbers of infiltrating immune cells are similar in NP-treated and control mice 2 days post-injury, fewer cells were present by the end of the first week post-injury. However, these results may be valid only for penetrating injuries rather than the more common contusion, which have differing immune cell dynamics and inflammation spread following injury, and therefore need to be verified in this model to before generalizing our results to all types of SCI.^{48,49} Additionally, mice were assigned a condition using a random number generator, so there may be human bias in assigning mice to treatment group. Further studies are

also needed to investigate direct effects of the NPs on trafficking of myeloid cell populations to the injury versus indirect effects on myeloid cell trafficking due to passive accumulation of NPs in the injury. In addition to altering the immune microenvironment in the spinal cord, NP administration leads to increased sparing of neurons. Early administration of NPs is able to substantially attenuate the early functional deficits. Administration of NPs at late time points is associated with early functional deficits, yet these functional deficits improve with time, with greater levels of improvement with earlier timing of NP administration. These NPs may provide a therapeutic option for patients with SCI to limit secondary damage, as they are formed from a material that has been approved by the FDA for use in other indications, and is stable and could be available off-the-shelf.

Declarations

Ethics approval and consent to participate:

All animal work was performed according to protocol number PRO00011456 approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Michigan.

Consent for publication:

Not applicable

Competing interests:

We declare a conflict of interest for Dr. Shea, who consults and has financial interests in Cour Pharmaceutical, which has licensed the nanoparticle technology and aims to commercialize the nanoparticles. Cour did not directly support the work submitted herein.

Data Availability

The datasets used and analyzed during the current study are available from the corresponding author upon reasonable request.

Acknowledgements

The authors would like to thank Jessica Chen for her mentorship, Rohit Maramraju for his cryosectioning help, and the UM Flow Cytometry Core for training and technical assistance. The graphical abstract was created using <https://BioRender.com>.

Funding

Funding was provided by the National Institutes of Health R01NS117103 and R01AI148076. S Schwartz has received funding from the National Institutes of Health T32TR004371.

Author Contributions

SEH and LDS designed the experiments. SEH completed all mouse work and surgeries. SEH primarily carried out the histology, imaging, qPCR, and flow cytometry experiments. BCR and

CRC helped with histology image analysis. BCR and SRS assisted with flow cytometry experiments. YW and CRC assisted with ladder beam analysis. SEH and LDS drafted the manuscript with revision help from BCR, SRS, BJC, and AJA. All authors approved the final manuscript.

References

1. Jain NB, Ayers GD, Peterson EN, et al. Traumatic spinal cord injury in the United States, 1993-2012. *Jama*. Jun 9 2015;313(22):2236-43. doi:10.1001/jama.2015.6250
2. Bowes AL, Yip PK. Modulating inflammatory cell responses to spinal cord injury: all in good time. *J Neurotrauma*. Nov 1 2014;31(21):1753-66. doi:10.1089/neu.2014.3429
3. Donnelly DJ, Popovich PG. Inflammation and its role in neuroprotection, axonal regeneration and functional recovery after spinal cord injury. *Exp Neurol*. Feb 2008;209(2):378-88. doi:10.1016/j.expneurol.2007.06.009
4. Gao TY, Huang FF, Xie YY, et al. Dynamic changes in the systemic immune responses of spinal cord injury model mice. *Neural Regen Res*. Feb 2021;16(2):382-387. doi:10.4103/1673-5374.290910
5. Liu Z, Yang Y, He L, et al. High-dose methylprednisolone for acute traumatic spinal cord injury: A meta-analysis. *Neurology*. Aug 27 2019;93(9):e841-e850. doi:10.1212/wnl.00000000000007998
6. Rouanet C, Reges D, Rocha E, Gagliardi V, Silva GS. Traumatic spinal cord injury: current concepts and treatment update. *Arq Neuropsiquiatr*. Jun 2017;75(6):387-393. doi:10.1590/0004-282x20170048
7. Sámano C, Nistri A. Mechanism of Neuroprotection Against Experimental Spinal Cord Injury by Riluzole or Methylprednisolone. *Neurochem Res*. Jan 2019;44(1):200-213. doi:10.1007/s11064-017-2459-6
8. Stirling DP, Liu S, Kubes P, Yong VW. Depletion of Ly6G/Gr-1 leukocytes after spinal cord injury in mice alters wound healing and worsens neurological outcome. *J Neurosci*. Jan 21 2009;29(3):753-64. doi:10.1523/jneurosci.4918-08.2009
9. Shechter R, London A, Varol C, et al. Infiltrating blood-derived macrophages are vital cells playing an anti-inflammatory role in recovery from spinal cord injury in mice. *PLoS Med*. Jul 2009;6(7):e1000113. doi:10.1371/journal.pmed.1000113
10. Kigerl KA, Gensel JC, Ankeny DP, Alexander JK, Donnelly DJ, Popovich PG. Identification of two distinct macrophage subsets with divergent effects causing either neurotoxicity or regeneration in the injured mouse spinal cord. *J Neurosci*. Oct 28 2009;29(43):13435-44. doi:10.1523/jneurosci.3257-09.2009
11. Chakraborty A, Ciciriello AJ, Dumont CM, Pearson RM. Nanoparticle-Based Delivery to Treat Spinal Cord Injury-a Mini-review. *AAPS PharmSciTech*. Mar 12 2021;22(3):101. doi:10.1208/s12249-021-01975-2
12. Yang Q, Lu D, Wu J, et al. Nanoparticles for the treatment of spinal cord injury. *Neural Regen Res*. Jun 1 2025;20(6):1665-1680. doi:10.4103/nrr.Nrr-d-23-01848
13. Cerqueira SR, Ayad NG, Lee JK. Neuroinflammation Treatment via Targeted Delivery of Nanoparticles. *Front Cell Neurosci*. 2020;14:576037. doi:10.3389/fncel.2020.576037
14. Mann AP, Scodeller P, Hussain S, et al. A peptide for targeted, systemic delivery of imaging and therapeutic compounds into acute brain injuries. *Nat Commun*. Jun 28 2016;7:11980. doi:10.1038/ncomms11980
15. Gu C, Geng X, Wu Y, et al. Engineered Macrophage Membrane-Coated Nanoparticles with Enhanced CCR2 Expression Promote Spinal Cord Injury Repair by Suppressing Neuroinflammation and Neuronal death. *Small*. Mar 2024;20(10):e2305659. doi:10.1002/smll.202305659


16. Yun X, Maximov VD, Yu J, Zhu H, Vertegel AA, Kindy MS. Nanoparticles for targeted delivery of antioxidant enzymes to the brain after cerebral ischemia and reperfusion injury. *J Cereb Blood Flow Metab.* Apr 2013;33(4):583-92. doi:10.1038/jcbfm.2012.209
17. Park J, Zhang Y, Saito E, et al. Intravascular innate immune cells reprogrammed via intravenous nanoparticles to promote functional recovery after spinal cord injury. *Proc Natl Acad Sci U S A.* Jul 23 2019;116(30):14947-14954. doi:10.1073/pnas.1820276116
18. Stirling DP, Yong VW. Dynamics of the inflammatory response after murine spinal cord injury revealed by flow cytometry. *J Neurosci Res.* Jul 2008;86(9):1944-58. doi:10.1002/jnr.21659
19. Carlson SL, Parrish ME, Springer JE, Doty K, Dossett L. Acute inflammatory response in spinal cord following impact injury. *Exp Neurol.* May 1998;151(1):77-88. doi:10.1006/exnr.1998.6785
20. Hellenbrand DJ, Quinn CM, Piper ZJ, Morehouse CN, Fixel JA, Hanna AS. Inflammation after spinal cord injury: a review of the critical timeline of signaling cues and cellular infiltration. *J Neuroinflammation.* Dec 7 2021;18(1):284. doi:10.1186/s12974-021-02337-2
21. Ciciriello AJ, Surnar B, Medy GD, Su X, Dhar S, Dumont CM. Biomaterial-targeted precision nanoparticle delivery to the injured spinal cord. *Acta Biomater.* Oct 15 2022;152:532-545. doi:10.1016/j.actbio.2022.08.077
22. Chen JY, Fu EJ, Patel PR, et al. Lentiviral Interleukin-10 Gene Therapy Preserves Fine Motor Circuitry and Function After a Cervical Spinal Cord Injury in Male and Female Mice. *Neurotherapeutics.* Jan 2021;18(1):503-514. doi:10.1007/s13311-020-00946-y
23. Smith DR, Dumont CM, Park J, et al. Polycistronic Delivery of IL-10 and NT-3 Promotes Oligodendrocyte Myelination and Functional Recovery in a Mouse Spinal Cord Injury Model. *Tissue Eng Part A.* Jun 2020;26(11-12):672-682. doi:10.1089/ten.TEA.2019.0321
24. Yang Y, De Laporte L, Zelivyanskaya ML, et al. Multiple channel bridges for spinal cord injury: cellular characterization of host response. *Tissue Eng Part A.* Nov 2009;15(11):3283-95. doi:10.1089/ten.TEA.2009.0081
25. Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A. CellProfiler 4: improvements in speed, utility and usability. *BMC Bioinformatics.* Sep 10 2021;22(1):433. doi:10.1186/s12859-021-04344-9
26. Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-image analysis. *Nat Methods.* Jun 28 2012;9(7):676-82. doi:10.1038/nmeth.2019
27. Berg S, Kutra D, Kroeger T, et al. ilastik: interactive machine learning for (bio)image analysis. *Nat Methods.* Dec 2019;16(12):1226-1232. doi:10.1038/s41592-019-0582-9
28. Saito E, Kuo R, Pearson RM, et al. Designing drug-free biodegradable nanoparticles to modulate inflammatory monocytes and neutrophils for ameliorating inflammation. *J Control Release.* Apr 28 2019;300:185-196. doi:10.1016/j.jconrel.2019.02.025
29. Carlson BM, Billington L, Faulkner J. Studies on the regenerative recovery of long-term denervated muscle in rats. *Restor Neurol Neurosci.* Jan 1 1996;10(2):77-84. doi:10.3233/rnn-1996-10203

30. Rowan SL, Rygiel K, Purves-Smith FM, Solbak NM, Turnbull DM, Hepple RT. Denervation causes fiber atrophy and myosin heavy chain co-expression in senescent skeletal muscle. *PLoS One*. 2012;7(1):e29082. doi:10.1371/journal.pone.0029082
31. Spector SA. Trophic effects on the contractile and histochemical properties of rat soleus muscle. *J Neurosci*. Aug 1985;5(8):2189-96. doi:10.1523/jneurosci.05-08-02189.1985
32. Wu P, Chawla A, Spinner RJ, et al. Key changes in denervated muscles and their impact on regeneration and reinnervation. *Neural Regen Res*. Oct 15 2014;9(20):1796-809. doi:10.4103/1673-5374.143424
33. Tosolini AP, Mohan R, Morris R. Targeting the full length of the motor end plate regions in the mouse forelimb increases the uptake of fluoro-gold into corresponding spinal cord motor neurons. *Front Neurol*. 2013;4:58. doi:10.3389/fneur.2013.00058
34. Fawcett JW, Curt A, Steeves JD, et al. Guidelines for the conduct of clinical trials for spinal cord injury as developed by the ICCP panel: spontaneous recovery after spinal cord injury and statistical power needed for therapeutic clinical trials. *Spinal Cord*. Mar 2007;45(3):190-205. doi:10.1038/sj.sc.3102007
35. Rhee P, Kuncir EJ, Johnson L, et al. Cervical spine injury is highly dependent on the mechanism of injury following blunt and penetrating assault. *J Trauma*. Nov 2006;61(5):1166-70. doi:10.1097/01.ta.0000188163.52226.97
36. Noble BT, Brennan FH, Popovich PG. The spleen as a neuroimmune interface after spinal cord injury. *J Neuroimmunol*. Aug 15 2018;321:1-11. doi:10.1016/j.jneuroim.2018.05.007
37. Milich LM, Ryan CB, Lee JK. The origin, fate, and contribution of macrophages to spinal cord injury pathology. *Acta Neuropathol*. May 2019;137(5):785-797. doi:10.1007/s00401-019-01992-3
38. Fleming JC, Norenberg MD, Ramsay DA, et al. The cellular inflammatory response in human spinal cords after injury. *Brain*. Dec 2006;129(Pt 12):3249-69. doi:10.1093/brain/awl296
39. Fehlings MG, Wilson JR, Tetreault LA, et al. A Clinical Practice Guideline for the Management of Patients With Acute Spinal Cord Injury: Recommendations on the Use of Methylprednisolone Sodium Succinate. *Global Spine J*. Sep 2017;7(3 Suppl):203s-211s. doi:10.1177/2192568217703085
40. Yin Z, Wan B, Gong G, Yin J. ROS: Executioner of regulating cell death in spinal cord injury. *Front Immunol*. 2024;15:1330678. doi:10.3389/fimmu.2024.1330678
41. Yu M, Wang Z, Wang D, Aierxi M, Ma Z, Wang Y. Oxidative stress following spinal cord injury: From molecular mechanisms to therapeutic targets. *J Neurosci Res*. Oct 2023;101(10):1538-1554. doi:10.1002/jnr.25221
42. Choi JI, Svensson CI, Koehn FJ, Bhuskute A, Sorkin LS. Peripheral inflammation induces tumor necrosis factor dependent AMPA receptor trafficking and Akt phosphorylation in spinal cord in addition to pain behavior. *Pain*. May 2010;149(2):243-253. doi:10.1016/j.pain.2010.02.008
43. Zhang L, Berta T, Xu ZZ, Liu T, Park JY, Ji RR. TNF- α contributes to spinal cord synaptic plasticity and inflammatory pain: distinct role of TNF receptor subtypes 1 and 2. *Pain*. Feb 2011;152(2):419-427. doi:10.1016/j.pain.2010.11.014

44. Li Y, Ritzel RM, Lei Z, et al. Sexual dimorphism in neurological function after SCI is associated with disrupted neuroinflammation in both injured spinal cord and brain. *Brain Behav Immun*. Mar 2022;101:1-22. doi:10.1016/j.bbi.2021.12.017
45. Stewart AN, Lowe JL, Glaser EP, et al. Acute inflammatory profiles differ with sex and age after spinal cord injury. *J Neuroinflammation*. May 13 2021;18(1):113. doi:10.1186/s12974-021-02161-8
46. Shi R, Kelly TM, Blight AR. Conduction block in acute and chronic spinal cord injury: different dose-response characteristics for reversal by 4-aminopyridine. *Exp Neurol*. Dec 1997;148(2):495-501. doi:10.1006/exnr.1997.6706
47. James ND, Bartus K, Grist J, Bennett DL, McMahon SB, Bradbury EJ. Conduction failure following spinal cord injury: functional and anatomical changes from acute to chronic stages. *J Neurosci*. Dec 14 2011;31(50):18543-55. doi:10.1523/jneurosci.4306-11.2011
48. Hausmann ON. Post-traumatic inflammation following spinal cord injury. *Spinal Cord*. Jul 2003;41(7):369-78. doi:10.1038/sj.sc.3101483
49. Beattie MS, Bresnahan JC. Cell Death, Repair, and Recovery of Function after Spinal Cord Contusion Injuries in Rats.

Graphical Abstract

After a C5 hemisection spinal cord injury, early intravenous administration of PLG nanoparticles reduces immune cell accumulation at the injury. This modulation of the inflammatory response is associated with increased neuron survival in the spinal cord, increased neuromuscular junction innervation in affected muscles, and improved motor performance.



Acute Timing Graphical Abstract (1).jpeg




Figure 1.jpg




Figure 2.jpg




Figure 3.jpg




Figure 4.jpg




Figure 5.jpg

After a C5 hemisection spinal cord injury, early intravenous administration of PLG nanoparticles reduces immune cell accumulation at the injury. This modulation of the inflammatory response is associated with increased neuron survival in the spinal cord, increased neuromuscular junction innervation in affected muscles, and improved motor performance.