Zuo Zhenjun (Orcid ID: 0000-0003-0263-4499)

Wang Zhong (Orcid ID: 0000-0003-1728-2932)

Author Manuscript

Title page

Title: Coordination between bioelements induce more stable macroelements than microelements in wetland plants

Running title: Coordination and stability of plant elementome

Author: Zhenjun Zuo[1,2], Peter B. Reich[3,4], Xiujuan Qiao[5], Haocun Zhao[1], Liangjian Zhang[1], Lei Yang[1], Tian Lv[1], Zhiyao Tang[2], Dan Yu[1*], Zhong Wang[1,6*]

Affiliations

[1] The National Field Station of Freshwater Ecosystems of Liangzi Lake, College of Life Sciences, Wuhan University, Wuhan 430072, China

[2] Institute of Ecology, College of Urban and Environmental Sciences, and Key Laboratory for Earth Surface Processes of the Ministry of Education, Peking University, Beijing 100871, China

[3] Institute for Global Change Biology, University of Michigan, Ann Arbor, Michigan 48109, USA

[4] Department of Forest Resources, University of Minnesota, St. Paul, Minnesota 55108 USA

[5] Key Laboratory of Aquatic Botany and Watershed Ecology, Wuhan Botanical Garden, Chinese Academy of Sciences [CAS], Wuhan 430074, China

[6] Key Laboratory of Biodiversity and Environment on the Qinghai-Tibetan Plateau, Ministry of Education, School of Ecology and Environment, Tibet University, Lhasa 850000, China

E-mail: Zhenjun Zuo, zuozhenjun@whu.edu.cn

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/ele.70025

This article is protected by copyright. All rights reserved.

Author Manuscript

Peter B. Reich, preich@umn.edu

Xiujuan Qiao, xjqiao@wbgcas.cn

Haocun Zhao, hczhao@whu.edu.cn

Liangjian Zhang, zhanglj@whu.edu.cn

Lei Yang, yangl@whu.edu.cn

Tian Lv, lvtian@whu.edu.cn

Zhiyao Tang, zytang@pku.edu.cn

Dan Yu, lakeyd@163.com

Zhong Wang, wangzhong@whu.edu.cn

*** Corresponding authors**

Zhong Wang, wangzhong@whu.edu.cn, ORCID: https://orcid.org/0000-0003-1728-2932

Dan Yu, lakeyd@163.com, ORCID: https://orcid.org/0000-0003-0954-8008

Address: 1. 299 Bayi Rd. Wuhan, 430072, China; 2. 10 Zangda East Rd. Lhasa, 850000, China

Tel: +86 27 68756834; **Fax**: +86 27 87882661

Keywords: ecological stoichiometry, evolutionary conservatism, functional biogeography, Stability of Well-coordinated Elements Hypothesis, Tibetan Plateau

Type of article: Letter

The number of words in Abstract: 155

The number of words in main text: 5137

The number of references: 59

The number of figures: 6

Author Manuscript

Author Contributions

Z. Wang designed research, Z. Zuo, H. Zhao, L. Zhang, L. Yang, T. Lv, and Z. Wang performed research and collected field data. Z. Zuo, H. Zhao, X. Qiao, and Z. Wang analyzed data. Z. Zuo and Z. Wang wrote the paper. P. B. Reich, X. Qiao, Z. Tang, and D. Yu helped interpret the data and revise the paper.

Data Availability Statement

The raw data and R code that supporting the findings of this study are available on Figshare at: https://doi.org/10.6084/m9.figshare.25952017.v4.

Author Manuscript

Coordination between bioelements induce more stable macroelements than microelements in wetland plants

ABSTRACT

Elements are the basic substances that make up living organisms, and the element composition in plants quantitatively reflect the adaptation of plants to environment. However, the drivers that constitute the species-specific plant elementome, as well the bivariate bioelemental correlations in determining the stability of different bioelements are yet unclear. Based on 1058 leaf observations of 84 plant species from 232 wetlands across large environmental gradients, we found that bioelements with higher concentration were more stable and evolutionary constrained. We proposed a Stability of Well-coordinated Elements Hypothesis, suggesting that bioelements that coordinate well in driving certain physiological functions constrain each other, thus maintaining relatively stable ratios in plants. In contrast, those functionally independent bioelements fluctuate greatly with environmental nutrient availability. Cold and saline stresses decreased plant stoichiometric networks connectivity, complexity and stability. Our research filled the gap in study of wetland plant elementome, and provided new evidences of plant-environment interactions in regions sensitive to climate change.

Author Manuscript

1 Introduction

All living beings are built with a specific range of the ratio of bioelements (Schimel, 2003; Sterner & Elser, 2002), namely, the elementome (Peñuelas et al., 2019). For plants, about 30 bioelements have been detected to serve vital but varied functions in participating in individual physiological processes and environmental adaptions (Ågren, 2008). Fundamentally, elementomes of organisms are determined by physiological requirements and are assumed to be species-specific and phylogenetically conserved (Atkinson et al., 2020; Sardans et al., 2021; Watanabe et al., 2007). However, the realized elemental composition is the consequence of multiple environmental filtering, including climate and habitat conditions (nutrient supply and stress, etc.) (Fernández-Martínez et al., 2021; Joswig et al., 2022; Zuo et al., 2022). In the classic framework of Hutchinson (1957)'s "*n*-dimensional hypervolume", the elementome is one of the best indicators to quantify the plant fitness to environments in multi-dimensions (Fernández-Martínez et al., 2021; Furey & Tilman, 2023; Peñuelas et al., 2019). Plants integrate multiple elements in perhaps optimal ratios to maximize their multifunctionality and to adapt to environmental heterogeneity (He et al., 2020), causing biogeographic patterns of the elementomes (Han et al., 2011; Reich & Oleksyn, 2004; Xing et al., 2021). Therefore, the variations in plant elemental composition along environmental gradients may reflect the joint controls of plant evolutionary history and present environments.

In terrestrial ecosystems, macroelements centered on C:N:P stoichiometry has been used to analyze plant-environment relationships (Han et al., 2011), identify species life-history strategies (Reich, 2014), probe community assembly mechanisms (Atkinson et al., 2020), determine the species biogeochemical niches (Fernández-Martínez et al., 2021; Peñuelas et al., 2019), and predict ecosystem responses to global change (Tang et al., 2018), whereas microelements also play important roles in homeostatic regulation, redox, and energy storage in plants (Kaspari, 2021; Lilay et al., 2024; Marschner, 2012). However, few studies have focused on the formation mechanisms of plant elementome *per se*, especially for both macroelements and microelements. In wetlands,

Author Manuscript

mineral elements carried by surface runoff from surrounding soils sink in sediments, producing more concentrated element micro-environments than upland soils (Herbert et al., 2015; Zuo et al., 2022). Since habitat elemental conditions in wetlands differ from those of terrestrial ecosystems, elemental stoichiometry of wetland plants is an ideal indicator reflecting the plant fitness to environments, as well as the generalities and differences in relation to environmental heterogeneity between wetland and non-wetland plants (Pan et al., 2020; Yang et al., 2021). However, no field study has linked plant elementome with plant phylogeny and functional biogeography in wetlands along large environmental gradients, especially under ongoing global change and hydrological regime changes (Pan et al., 2023).

Wetlands are distinguished from upland habitats with unique biogeochemical processes, light, CO_2 and O_2 availability and hydrological regime (Moor et al., 2017). Wetland plants are characterized by sufficient water availability and nutrient supply in sediments, but suffer from additional stresses of osmotic pressure, gas limitation, and strong hydraulic forces in water (Maberly & Gontero 2018), and thus evolved a suite of morphological and eco-physiological traits to adapt to water saturated environments (*e.g.*, leaf gas film formation and enhanced shoot and root porosity, Pan et al., 2023). In the economic strategy dimension, the leaf economics spectrum (LES), which was built based on coordination among six mass-based leaf traits, indicates that plants with high leaf N and P concentrations and specific leaf area show quick-return strategies on investment of nutrients (Reich 2014; Reich et al., 1997; Wright et al., 2004). Among the plant growth forms, wetland plants are ones often using quick-return strategies (Pan et al., 2020; Yang et al., 2021). Previous studies have focused on plant photosynthetic physiology (Pedersen et al., 2013), and leaf C:N:P stoichiometry (Hu et al., 2021; Wang et al., 2015) in wetland habitats. However, as the

Author Manuscript

essential participants in plant physiological processes, the stoichiometry of bioelements other than C, N and P in wetland plants and their controlling factors are yet unclear.

The alpine and arid regions in western China are often considered as amplifiers for climate change with continued warming, spatially heterogeneous changes in precipitation and anthropogenic disturbances (Piao et al., 2010; Song et al., 2021), resulting in the loss or salinization of wetlands. Wetland plants in these regions are exposed to harsh environments such as low temperature and high salinity (Wang et al., 2015), shaping its realized elementome from the phylogeny-determined fundamental elementome by both climate and habitat filtering (Figure 1a). However, the shaping processes are bioelement-specific, and lead to differences in bioelemental stability. Furthermore, the realized elementome is also the consequence of coordination among bioelements, but received less attention. Environmental change not only affect the bioelemental concentrations, but adjust the complex networks among bioelements (He et al., 2020; Li et al., 2022), then forming the plant elementome (Figure 1b). Based on 1058 leaf samples collected by uniform protocols in the study regions (Figure S1), we aimed to clarify the stoichiometric patterns and their phylogenetic and environmental controls of plant elementome, and explore the variations of plant element networks in stressful conditions. We hypothesized that environmental changes induce allometric accumulations of different bioelements, alter the connectivity and complexity of the plant element networks, but plant bioelements that coordinate well with each other maintain higher stability in different environments, and *vice versa*.

2. Methods and Materials

2.1 Study region and field investigation

This study region ranges from 27.4°N to 48.7°N, 76.2°E to 109.8°E and 195m to 5127m a.s.l., and covers almost all the alpine and arid regions in western China (Figure S1). This region is characterized by alpine and arid climate, with mean annual air temperature and mean annual

Author Manuscript

precipitation ranging from −4.7°C to 12.5°C and from 17.2mm to 685.4mm, respectively (Song et al., 2021).

The alpine and arid regions of western China are rich in various wetland habitats (Wang et al., 2015; Zuo et al., 2022), yet also face the impacts of climate change and anthropogenic activities, which can alter the nutrient resources and water resources of wetland ecosystems (Piao et al., 2010). The habitats we investigated included shallow areas (no deeper than 1.5 m) of lakes, rivers, and marshes, as well as streams, ponds, and channels, while ephemeral water bodies were excluded. Moreover, we recorded the information of sampling site as summarized in Table S1.

We conducted a large-scale field investigation across the western China during the growing season in 2018 and 2019. Detailed methods of field sampling and laboratory analysis were provided in Appendix S1. Briefly, sediment, water and plant (including emergent, floating-leaved and submerged plants) samples were collected in 232 wetlands, and 17 elements (C, N, P, S, K, Ca, Mg, Na, Fe, Al, Mn, Si, B, Zn, Cu, Ni and Co) of the sediment (232 samples), water (232 samples) and plants (1058 leaf samples) were measured.

2.2 Data analysis

2.2.1 Data arrangement

For each of the 17 bioelements, arithmetic mean value, standard deviation, maximum and minimum values, and coefficient of variation (CV) for overall species, emergent plants, floating-leaved plants, and submerged plants were sorted out respectively. The CV, which was calculated by dividing standard deviation by mean value, was introduced to represent the variability in bioelemental concentrations (Han et al., 2011; Karimi & Folt 2006). Marschner (2012) provided the average concentrations of 14 mineral elements in plant shoot dry matter that are sufficient for adequate growth, including 12 elements (Ca, K, N, S, Mg, P, Fe, Mn, B, Zn, Cu and Ni) in this

Author Manuscript

study. The concentration that is not limit plant growth are considered as the physiological requirement of a certain bioelements in plants.

Before analysis, all data of bioelemental concentrations were checked for normality of residual using Shapiro-Wilk normality tests and homogeneity of variances by Bartlett tests. Then the non-parametric methods (Kruskal-Wallis test and Dunn's Post Hoc Multiple Comparisons) were used to compare the differences in bioelemental concentrations among life-forms.

In this study, growing season mean air temperature (GST) is the mean temperature from May to September, while growing season precipitation (GSP) is the total precipitation of the five months (Wang et al., 2015). We used eight environmental variables, including two climatic variables (GST and GSP), three sediment variables (elemental concentration, pH and electrical conductivity) and three water variables (elemental concentration, pH and salinity). Each bioelement was fitted against the environmental variables, as well as latitude and altitude, by simple linear regressions to illustrate the bioelement-environment correlations.

2.2.2 Linear mixed effect model

We constructed a phylogenetic tree of 84 sampled species to test the phylogenetic signals (Blomberg's *K*) of 17 bioelements of wetland plants (Blomberg et al., 2003). Detailed methods of phylogenetic signal detection are provided in Appendix S2.

The linear mixed effect model (LMM) and variance partitioning analysis were employed to partition the bioelement variations into phylogeny and environments by using residual maximum likelihood estimation (Watanabe et al., 2007). We excluded plant families from which less than one species was sampled (Zhao et al., 2016). The phylogenetic effects were represented by a hierarchically nested structure "order/family/species". This method enabled us to investigate the degree of variation at each phylogenetic level. Variations in bioelemental concentrations effected by environmental variables (including climate, sediment, and water properties) were assigned to the 'site' component of the model (Watanabe et al., 2007; Zhao et al., 2016). The overall random

term within LMM was written as 'site + [(order/family/species)]', and no fixed factor was defined. We used the "*lmer*" function of the "*lme4*" package in *R* (Bates et al., 2015).

2.2.3 Phylogenetically controlled linear mixed model

We tested the effects of climate, sediment, and water properties on elementome and stoichiometric ratios between macroelements of wetland plants using Bayesian phylogenetic linear mixed models and the R "*brms*" package (Bürkner, 2017). In each Bayesian model, we used eight predictor variables (GST, GSP, sediment pH, sediment electrical conductivity, water pH, water salinity, and concentrations of corresponding elements in sediment and water) as fixed effects, with species as a random effect and controlled for phylogenetic effects using phylogenetic tree (Fernández-Martínez et al., 2021; Sardans et al., 2021). Priors used followed a normal distribution (0, 2) for beta estimates of the predictors and a *t*-student distribution (3, 0, 5) for the intercept, random and phylogenetic effects, and the residuals (Fernández-Martínez et al., 2021). Based on the null hypothesis of Mean = 0, we further estimated the probability of direction and Bayes factors of all predictors using the R "*bayestestR*" package (Makowski et al., 2019), as an estimation of the relevance of predictor variables.

2.2.4 Partial least squares path modeling (PLS-PM)

To explore the underlying regulatory mechanisms of environmental patterns of bioelements, partial least squares path modeling was further applied to identified the direct and indirect effects of geography, climate, sediment, and water properties on variations in bioelements. We performed PLS-PM for eight macroelements (C, N, P, S, K, Ca, Mg and Na) and nine microelements (Fe, Al, Mn, Si, B, Zn, Cu, Ni and Co), respectively. All the 11 observation variables were classified into four latent variables – geography (latitude, longitude and altitude), climate (GST and GSP), sediment properties (sediment pH, electrical conductivity and elemental concentrations), and water properties (water pH, salinity, and elemental concentrations). The plant bioelements were controlled by all the observation variables directly or indirectly. The models were constructed by

Author Manuscript

Author Manuscript

"*innerplot*" function of the "*plspm*" package in *R* (Sanchez 2013). Goodness of fit index (*GoF*), R^2, path coefficients (standardized direct effects) and standardized indirect effects were used to estimate model performance and compare the effects of different latent variables on the patterns of elements.

2.2.5 Leaf element network (LEN) analysis

With reference to plant trait network (He et al., 2020), we introduced LEN to clarify the complex relationships between bioelements and further assess the variations in the synergies of bioelements across environmental gradients. Detailed methods of LEN analysis are provided in Appendix S3 and Table S2. Briefly, we selected temperature and water salinity as proxies for climate and salt stresses, and constructed LENs at different temperature and salinity levels. We employed six overall parameters (edge density, average path length, diameter, average clustering coefficient, modularity, and vulnerability) to quantify the "connectivity", "complexity" and "stability" of LENs, and five node parameters (degree, weighted degree, closeness, betweenness and clustering coefficient) to assess the "connectedness" and "centrality" of bioelements within LENs (Felipe-Lucia et al., 2020, He et al., 2020). We further tested the stability of the node parameters for 17 bioelements in different conditions, to explore the consistency of the importance of each bioelement across environmental gradients (Li et al., 2022). If a certain bioelement with high node parameter shows high stability, it means that such bioelements always play central roles in different environmental conditions (Rao et al., 2023).

3 Results

3.1 Stoichiometric characteristics of wetland plants

For all the 1058 observations, the mean concentrations of the 17 bioelements varied by six orders of magnitude from the most (C: 374.93 mg g^{-1} or 31.25 mol kg^{-1}) to the least abundant one (Co: 0.0024 mg g^{-1} or 0.41×10^{-4} mol kg^{-1}, Figure 2a, Table S3), with mass-based ratio of

C_{155305}:Ca_{12747}:K_{10186}:N_{9822}:Na_{3706}:S_{2631}:Mg_{2621}:P_{1191}:Fe_{1152}:Al_{701}:Mn_{518}:Si_{176}:$B_{64.8}$:$Zn_{26.7}$:$Cu_{2.99}$:$Ni_{2.09}$:Co_1 or atom ratio of C_{763580}:Ca_{18802}:K_{15409}:N_{41392}:Na_{9508}:S_{4852}:Mg_{6445}:P_{2267}:Fe_{1214}:Al_{1531}:Mn_{555}:Si_{370}:B_{347}:$Zn_{24.2}$:$Cu_{2.76}$:$Ni_{2.12}$:Co_1. Considering the high concentrations of Na, we defined it as a macroelement in this study. Among the three life-forms, there were different bioelemental concentrations except for S (Figure 2a). Submerged plants concentrated more mineral elements, but had less leaf C and K than floating-leaved plants and emergent plants. Both N and P showed the highest concentrations in floating-leaved plants.

The variations of the 17 bioelements differed greatly. The nine microelements varied more (CV = 87%-188%) than the eight macroelements (CV = 13%-85%, Figure 2b). The CV was negatively correlated with both the mean concentrations of bioelements (R^2 = 0.36, P = 0.01, Figure 2c) and their physiological requirements (R^2 = 0.33, P < 0.05, Figure 2d), indicating that the bioelements with higher concentrations and higher requirements were more stable in plants.

3.2 Geographic and environmental patterns of bioelements

For all the 17 bioelements of overall 1058 observations, four (Na, S, Mg and Zn) increased, whereas seven (K, P, Fe, Al, Si, B and Co) decreased towards high latitude (Figure S3a, S4). Because of the special topographical features in the study areas, the high-altitude Tibetan Plateau lies in the south, altitudinal trends of bioelements were opposite to latitudinal gradients (Figure S5). Sixteen of the 17 bioelements (except Ca) were significantly correlated with GST, of which, Na, S, Mg, Mn, Zn, Cu and Ni increased while C, K, N, P, Fe, Al, Si, B and Co decreased with increasing GST (Figure S3c, S6). The bioelement-GSP relationships were opposite to the bioelement-GST relationship (Figure S7).

All of the 17 bioelements showed significant positive correlations with their corresponding nutrient levels in sediments (Figure S3e, S8). Fourteen bioelements positively, but K negatively correlated with their water nutrients, while P showed no correlations (Figure S9). Six bioelements

Author Manuscript

Author Manuscript

(Ca, Na, S, Mg, Si and B) increased, whereas 10 (K, N, P, Fe, Al, Mn, Zn, Cu, Ni and Co) decreased towards high sediment or water pH (Figure S10, S11). With increasing environmental stress towards high water salinity, concentrations of Na, S, Mg and B increased, but all the other 13 bioelements decreased (Figure S3i, S12, S13). Furthermore, these biogeographic patterns of the 16 bioelements were basically consistent among three life-forms (Table S4).

3.3 Phylogenetic and environmental controls on bioelements

Blomberg's *K* statistical identified 15 of 17 bioelements with significant phylogenetic signals (except Mg and B), with *K* values ranged from 0.089 (S and Cu) to 0.223 (K) (Table S5). LMM revealed that phylogeny and environments co-regulated the variations in bioelemental concentrations (Figure 3a), with a mean degree of explanation 23% (with ranges of 4%-56%) by phylogenetic level (incorporating effects of order, family and species levels) and 36% (7%-56%) by sites (Table S6). However, phylogenetic level accounted for a greater amount of the variation in macroelements (31%) than microelements (15%). On the contrary, sites explained microelements (44%) more than macroelements (27%). Furthermore, the variance explained by phylogeny positively correlated with bioelemental concentrations at all phylogenetic levels ($P < 0.05$; Figure 3b), while those by environments (sites) increased with decreasing bioelemental concentrations ($P < 0.001$; Figure 3c). That is, the higher concentrations of bioelements, the more regulated by phylogeny, and the lower concentrations of bioelements, the more regulated by environments.

Bayesian phylogenetic linear mixed models explained 28% (N)-64% (Na) of the variations in bioelements (Figure 4a, Table S7). Within environmental variables, nutrient supplies in sediment and water affected the variations in most bioelements, especially for microelements. Climate, especially GST, explained more for macroelements than microelements (Figure 4b). Correspondingly, the models explained 24% (N:P)-61% (C:Ca) of the variations in 28 stoichiometric ratios (Figure S14). Sediment elemental concentration and GST were the main

drivers of the variations in most stoichiometric ratios, and water salinity also showed high correlations with Na- and S-relevant stoichiometric ratios.

PLS-PM explained 47% and 43% of the variances of macroelements and microelements, but illustrated different regulatory mechanisms for environmental patterns of the two bioelement groups (Figure 4c-f, Table S8). For macroelements, water properties, climate and sediment properties positively affected the bioelements significantly with the standardized direct effects of 0.37, 0.24 and 0.24 (Figure 4c), but climate presented to be the strongest explanatory variable with considering the total effect of 0.65 (Figure 4d). For microelements, sediment properties (0.34) and water properties (0.30) had significantly positive effects, but climate showed no direct effects ($P >$ 0.05, Figure 4e). Sediment properties presented to be the first explanatory variable with the total effect of 0.53 (Figure 4f). Geographic variables indirectly, via influencing climate, sediment, and water properties, affected both the macroelements and microelements.

3.4 Variations in leaf element networks (LENs) along GST and salinity gradients

Most of the 17 bioelements showed significant correlations between each other (Figure 5a). Leaf C weakly positively correlated with N, P and K, and more strongly negatively correlated with all the other bioelements (except Zn with R= -0.05 and P = 0.096). N, P and six metallic microelements (Fe, Al, Mn, Zn, Cu, Ni) showed positive correlations with each other (Figure S15). Towards stressful conditions (low temperature and high salinity), the edge density and average path length of LENs decreased, whereas the diameter, average clustering coefficient and vulnerability increased (Figure 5b). Modularity was higher at low-GST but lower at saline water (Figure 5b). Variations in overall parameters of networks showed that the connectivity, complexity, and stability of LENs were lower in stressful conditions, and thus indicated that the synergies of bioelements became weaker. Euclidean distances between paired LENs showed that, both along temperature

Author Manuscript

Author Manuscript

and salinity gradients, the greater the environmental differences, the more disparities of LENs (Appendix S4, Figure S16).

For node parameters of LENs, leaf C was the hub trait with high degree at all temperature levels and the lower three salinity levels (Figure S17), while the degree of some stress resistance related bioelements (*e.g.*, S) increased sharply at the highest salinity level (Figure S17b). Environmental stress rearranged the module composition (Figure S18). Additionally, the stability of the five node parameters (degree, weighted degree, closeness, betweenness and clustering coefficient) for the 17 bioelements positively correlated to the corresponding parameter values (Figure S19), indicating that the important bioelements (*e.g.*, hub traits, C) tended to maintain their importance across environmental gradients.

4 Discussion

4.1 Differentiation in elemental stoichiometry among life-forms

Leaf bioelements varied significantly among different plant life-forms in wetlands. Submerged plants possessed less C than other life-forms (Hu et al., 2021; Wang et al., 2015). Physiologically, C in plants is mainly invested in carbon skeletons as structural C (C_S), for example in the cell wall, to support plants (Ma et al., 2018). However, the additional support from water buoyancy reduces the needed strength of plant supporting tissues, and thus lowers the requirement for C, which always is a limiting element for plant photosynthesis underwater (Maberly & Gontero, 2018; Pedersen et al., 2013). Further calculation about C_S concentrations in this study demonstrated that both the absolute values and relative amount of the C_S in submerged plants were significantly lower than that in the other two life-forms of plants (Appendix S5, Table S3).

For the mineral elements, 13 of the 16 bioelements were detected in the highest concentrations in submerged plants (Figure 2a), reflecting the special elementome of this life-form in adapting to the submerged environment. For example, Ca-containing calcium pectate (Xing et al., 2021) and

Author Manuscript

Si (Schoelynck & Struyf, 2016) serve load-bearing functions in cell walls, enhancing the flexibility and resistibility of submerged plants to hydraulic forces. In underwater environments, plants suffer from reduced light penetration and CO_2 diffusion, which constrain the photosynthesis of submerged plants (Pedersen et al., 2013). Considering that the light-driven photosynthesis is enzymatic-dependent process, the enrichment of ionic elements, such as Mg, Mn, Fe, Cu (Ågren, 2008; Grusak et al., 2016), for the synthesis of chlorophyll, enzymes (*e.g.*, Rubisco) and coenzyme factors in plants help to improve the limited photosynthetic potential underwater. Unlike emergent plants, submerged leaves are surrounded by water and thus directly face the osmotic stress from the solute in water. Submerged plants reserve more inorganic (*e.g.*, Na^+, SO_4^{2-}) or low molecular weight organic compounds (*e.g.*, S-containing compounds) as osmoprotectants in leaves to maintain osmotic balance (Deinlein et al., 2014; Zuo et al., 2022). In this study, the higher the element contents in water, the more enriched the bioelements in submerged plant leaves (Figure S9). Furthermore, the alkaline microenvironment around leaves produced by photosynthesis may promote the enrichment of metal cations in leaves (Pedersen et al., 2013). In addition, lower C concentrations is also a reason inducing higher mass-based fractions of the mineral elements in submerged plants.

4.2 Variability of bioelements and the relative effects of phylogeny and environments

In this study, we extracted three highlights illustrating the regulatory mechanisms of variations in wetland plant bioelements in alpine and arid regions (H1-H3 in Figure 6a,b).

Firstly, wetland plants were more stable in macroelements than microelements, that is, the higher the concentrations, or the higher the physiologically requirement, the more stable (smaller CV) the bioelements are in plants (Figure 2c,d), coinciding with the Stability of Limiting Elements Hypothesis (Han et al., 2011). The differences in stability between macroelements and microelements involves the degree to which each bioelement once acquired will cause a shift in the

Author Manuscript

concentrations of other bioelements, which for example are required in certain ratios to build certain tissues or to drive specific processes and functions. Macroelements with high demands (N, P, K, *etc*.) are always the key elements limiting productivity (*i.e.* accumulation of C at the whole plant scale). In this study, N, P and K functionally coordinated to each other tightly (Figure 5a). A change in one macroelement will constrain or influence the shift in others. For example, more N accumulation might lead to more P and K absorption, and promote C fixation by enhancing photosynthesis. More C accumulation create the "nutrient dilution effect" (Kaspari, 2021), which in turn reduces the mass-based N concentration and keeps it relative stable. In contrast, microelements had weak relationships to macroelements, even negatively correlated to C (Figure S15), showing relatively independent function in driving specific processes. In addition, the microelements are almost always found at levels much beyond where they influence function (Kaspari, 2021). Thus, if there is more available in environment, the microelements passively end up with more (*e.g.* what is sometimes called luxury absorption) but it doesn't change N or C concentrations. Similarly, if the microelements are passively acquired, it doesn't matter what the N and C concentrations are. Hence, the microelements vary more in lockstep with availability. In summary, we proposed a Stability of Well-coordinated Elements Hypothesis, suggesting that bioelements which coordinate together well in driving certain physiological functions (*e.g.* biomass production) constrain each other and thus are kept relatively stable in plants, while those functionally independent bioelements fluctuated greatly with the nutrient availability in environment.

Secondly, macroelements are more evolutionary constrained, while microelements are more susceptible to environmental influences. Both the physiological requirements and environmental nutrient limitations can influence the regulation strength of phylogeny and environments on bioelement variations (Ågren, 2008; Peñuelas et al., 2019). Macroelements are components of biological macromolecules, often in fixed ratios and coordinated relations, and thus their strong

Author Manuscript

stabilities perhaps also contribute to the evolutionary conservatism of the macro-elementome to maintain in different taxa (Wang et al., 2015; Zhao et al., 2016). In contrast, most microelements are the main components of coenzymes (Grusak et al., 2016), which are functionally independent to macroelements and more sensitive to environmental nutrients (Han et al., 2011). Fundamental biochemical processes have distinct rates across species, which induce different requirements for various bioelements, resulting in different regulations of macroelements and microelements (Sardans et al., 2021). To date, there have been few studies focusing on the phylogenetic signals of plant elementome, especially microelements (but see Fernández-Martínez et al., 2021). Evolutionary history (Ackerly, 2009), genetic variation (Crisp & Cook, 2012) and stable selection (Donoghue, 2008) could control the variations in species-specific plant functional traits. In this study, the relative effects of phylogeny on bioelements increased while that of environments decreased with increasing bioelemental concentrations (Figure 3).

Thirdly, environmental variation had greater effects on microelements than macroelements, which may be because microelements are environment sensitive, physiologically required in low amounts, and have weak functional coordination with other bioelements (Han et al., 2011; Karimi & Folt, 2006; Zhao et al., 2016). Within environmental factors, nutrient supplies are the primary ones in determining the patterns of microelements, but nutrient supplies, stress factors and climate control the variations in macroelements jointly. Variations in macroelements are influenced by climate (Han et al., 2011; Reich & Oleksyn, 2004), soil factors (Joswig et al., 2022), water chemistry (Fernández-Martínez et al., 2021) and environmental stresses (*e.g.*, salinity, Zuo et al., 2022). In this study, climate had significant direct and indirect effects on macroelements, while only a weak indirect effect on microelements (Figure 4c-f). Temperature and precipitation are critical factors determining plant growth and survival in alpine and arid regions (Shen et al., 2022). Directly, climatic factors regulate the physiological metabolic process and growth rate of plants, and thus affect the absorption and accumulation of elements (the Temperature-Plant Physiological

Author Manuscript

Hypothesis, Reich & Oleksyn, 2004). Indirectly, climatic factors regulate microbial mineralization and biogeochemical cycles of elements (Delgado-Baquerizo et al., 2013), resulting in heterogenetic resource availability and environmental stresses in different environments, further causing nutrient limitation of macroelements (*e.g.*, N and P in the Temperature-Biogeochemical Hypothesis, Reich & Oleksyn, 2004). However, microelements are mainly determined by the nutrient status of soil parent material rather than climatic factors, and thus are not the limiting elements in environments, which are consistent with our results (Figure 4). The specific biological function, weak mobility and high environmental plasticity of microelements together lead to their high dependence on nutrient supply.

This study further emphasized the significant effects of environmental nutrients on bioelements beyond C, N and P, especially microelements, which had rarely been mentioned in previous studies. Our findings highlighted that intrinsic species identities, theoretical physiological requirements and practical environmental constraints jointly optimize the elementome of plants (Peñuelas et al., 2019). However, further studies on the homeostasis/plasticity of bioelements would be much helpful for understanding the linkages between plant elementome and species adaptation and ecosystem functioning (Fernández-Martínez, 2022).

4.3 Linking element network with plant response to environment

Beyond concentrations, correlations among bioelements and their variations also reflect the plant response to environments (He et al., 2020). In this study, the correlations among bioelements showed lower connectivity, complexity, and stability in cold and saline habitats (Figure 5b), indicating that stressful environments restricted inter-element synergism in plants (Li et al., 2022; Rao et al., 2023). Maintaining stable and complex bioelemental correlations requires high construction cost, and may be disadvantageous for plants in stressful habitats (Alon, 2003; Flores-Moreno et al., 2019). In contrast, networks with high connectivity, complexity and stability in non-

Author Manuscript

stressed environments may signify a tight integration among bioelements which allows plants to access resources more efficiently (Rao et al., 2023), namely, "complexity begets stability" (Yuan et al., 2021).

Plants responded to cold and saline environments by adjusting the correlations among the bioelements, which could be detected from the variations in modular composition and node parameters of LENs. Firstly, changes in network connectivity have been considered as the trade-off between connection efficiency and connection cost (Flores-Moreno et al., 2019). LENs exhibited lower connection (*e.g.*, lower edge density) among elements to reduce the cost of bioelemental collaboration under stressful environments. In addition, we found a higher network modularity at low-GST level (Figure 5b), which allowed plants to perform their specific functions more independently, and prevented one module failure from spreading to other modules (Gilarranz et al., 2017; Rao et al., 2023). Secondly, maintaining the robust correlations among certain bioelements is an important basis for plants to accomplish physio-biochemical processes. N and P always constituted a module across temperature and salinity gradients (Figure 5a), which was consistent with the results of woody plants across climatic regions (Flores-Moreno et al., 2019). N and P are central to the leaf economic spectrum and their robust correlation is critical to the resource trade-off strategies of plants in diverse environments (Reich, 2014; Reich et al., 2010; Reich & Oleksyn, 2004; Wright et al., 2004). Similarly, Al, Fe, Ni and Co always constituted a module to maintain necessary biochemical reactions (Grusak et al., 2016). Thirdly, LENs could shift the connectedness and centrality of bioelements to respond to stresses. For example, the hub traits, C, consistently maintained their high importance at all levels of environments (Figure S19), and exhibited the highest stability across environmental gradients in this study, indicating the important role of C on plant phenotype and biomass production (Ma et al., 2018). On the contrary, the degree of S increased with increasing salinity, confirming the crucial role of S on salt resistance (Zuo et al., 2022). Therefore, we inferred that connections related to hub traits are more robust, while

Author Manuscript

correlations among non-hub traits are more likely to change or even break (*e.g.*, the decoupling of plant N-S correlation in high-salt environment, Zuo et al., 2022), resulting in deformations of LENs with environmental changes (Figure 6c).

5. Conclusion

Using multidimensional approaches, we evaluated the variations in wetland plant elementome and their correlations beyond C, N and P across life-forms, phylogeny, climate, and environments in alpine and arid regions. The variation patterns of bioelements suggest general rules about how phylogeny and environments co-regulate the elementome. Bioelements with higher concentrations were more stable, and the drivers of the variations in bioelements shifted from environmental modification to evolutionary constraint. Tighter correlations among bioelements might mean better coordination in driving physiological functions, which are regulated by species-specific requirements, and constrained by functional requirements for certain ratios, resulting in less variable concentrations. In contrast, functionally independent bioelements with weak coordination are more influenced by nutrient availability in habitats. The variations of microelements were primarily controlled by nutrient supplies, while macroelements were mainly affected by climate, followed by environmental nutrients and stresses. Stressful (cold or saline) habitats lead to looser leaf element correlations and lower inter-element coordination and stabilities. The highlights extracted in this study advance our understanding of the causes and consequences of wetland plant elementome in harsh environments.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (Grant 32360287 by Zhong Wang), the Special Project of Basic Work of Science and Technology,

Author Manuscript

Ministry of Science and Technology, China (Grant Number 2013FY112300 by Dan Yu) and the ASCEND Biological Integration Institutes grant NSF-DBI-2021898 by Peter B. Reich. We thank three anonymous reviewers and the editor for their constructive comments and suggestions to improve this manuscript.

Conflicts of Interest

The authors declare no competing interests.

Supporting Information

Additional supporting information can be found online in the Supporting Information section at the end of this article.

References

Ackerly, D. (2009) Conservatism and diversification of plant functional traits: evolutionary rates versus phylogenetic signal. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 19699–19706.

Ågren, G.I. (2008) Stoichiometry and nutrition of plant growth in natural communities. *Annual Review of Ecology, Evolution, and Systematics*, 39, 153–170.

Alon, U. (2003) Biological networks: the tinkerer as an engineer. *Science*, 301, 1866–1867.

Atkinson, C.L., van Ee, B.C. & Pfeiffer, J.M. (2020) Evolutionary history drives aspects of stoichiometric niche variation and functional effects within a guild. *Ecology*, 101, e03100.

Bates, D., Maechler, M., Bolker, B.M. & Walker, S.C. (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67, 1–48.

Blomberg, S.P., Garland, T. & Ives, A.R. (2003) Testing for phylogenetic signal in comparative data:

Author Manuscript

behavioral traits are more labile. *Evolution*, 57, 717–745.

Büerkner, P.-C. (2017) brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, 80, 1–28.

Crisp, M.D. & Cook, L.G. (2012) Phylogenetic niche conservatism: what are the underlying evolutionary and ecological causes? *New phytologist*, 196, 681–694.

Deinlein, U., Stephan, A.B., Horie, T., Luo, W., Xu, G. & Schroeder, J.I. (2014) Plant salt-tolerance mechanisms. *Trends in Plant Science*, 19, 371–379.

Delgado-Baquerizo, M., Maestre, F.T., Gallardol, A., Bowker, M.A., Wallenstein, M.D., Luis Quero, J. et al. (2013) Decoupling of soil nutrient cycles as a function of aridity in global drylands. *Nature*, 502, 672–676.

Donoghue, M.J. (2008) A phylogenetic perspective on the distribution of plant diversity. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 11549–11555.

Felipe-Lucia, M.R., Soliveres, S., Penone, C., Fischer, M., Ammer, C., Boch, S. et al. (2020) Land-use intensity alters networks between biodiversity, ecosystem functions, and services. *Proceedings of the National Academy of Sciences of the United States of America*, 117, 28140–28149.

Fernández-Martínez, M. (2022) From atoms to ecosystems: elementome diversity meets ecosystem functioning. *New Phytologist*, 234, 35–42.

Fernández-Martínez, M., Preece, C., Corbera, J., Cano, O., Garcia-Porta, J., Sardans, J. et al. (2021) Bryophyte C:N:P stoichiometry, biogeochemical niches and elementome plasticity driven by environment and coexistence. *Ecology Letters*, 24, 1375–1386.

Flores-Moreno, H., Fazayeli, F., Banerjee, A., Datta, A., Kattge, J., Butler, E.E. et al. (2019) Robustness of trait connections across environmental gradients and growth forms. *Global Ecology and Biogeography*, 28, 1806–1826.

Author Manuscript

Furey, G.N. & Tilman, D. (2023) Plant chemical traits define functional and phylogenetic axes of plant biodiversity. *Ecology Letters*, 26, 1394–1406.

Gilarranz, L.J., Rayfield, B., Linan-Cembrano, G., Bascompte, J. & Gonzalez, A. (2017) Effects of network modularity on the spread of perturbation impact in experimental metapopulations. *Science*, 357, 199–201.

Grusak, M.A., Broadley, M.R. & White, P.J. (2016) Plant macro- and micronutrient minerals. *eLS*, 1–6.

Han, W.X., Fang, J.Y., Reich, P.B., Woodward, F.I. & Wang, Z.H. (2011) Biogeography and variability of eleven mineral elements in plant leaves across gradients of climate, soil and plant functional type in China. *Ecology Letters*, 14, 788–796.

He, N., Li, Y., Liu, C., Xu, L., Li, M., Zhang, J. et al. (2020) Plant trait networks: improved resolution of the dimensionality of adaptation. *Trends in Ecology & Evolution*, 35, 908–918.

Herbert, E.R., Boon, P., Burgin, A.J., Neubauer, S.C., Franklin, R.B., Ardon, M. et al. (2015) A global perspective on wetland salinization: ecological consequences of a growing threat to freshwater wetlands. *Ecosphere*, 6, 1–43.

Hu, J., Yu, H., Li, Y., Wang, J., Lv, T., Liu, C. et al. (2021) Variation in resource allocation strategies and environmental driving factors for different life-forms of aquatic plants in cold temperate zones. *Journal of Ecology*, 109, 3046–3059.

Hutchinson, G.E. (1957) Concluding remarks. *Cold Spring Harbor Symposia on Quantitative Biology*, 22, 415–427.

Joswig, J.S., Wirth, C., Schuman, M.C., Kattge, J., Reu, B., Wright, I.J. et al. (2022) Climatic and soil factors explain the two-dimensional spectrum of global plant trait variation. *Nature Ecology & Evolution*, 6, 36–50.

Karimi, R. & Folt, C.L. (2006) Beyond macronutrients: element variability and multielement

Author Manuscript

stoichiometry in freshwater invertebrates. *Ecology Letters*, 9, 1273–1283.

Kaspari, M. (2021) The invisible hand of the periodic table: how micronutrients shape ecology. *Annual Review of Ecology, Evolution, and Systematics*, 52, 199–219.

Li, Y., Liu, C., Sack, L., Xu, L., Li, M., Zhang, J. et al. (2022) Leaf trait network architecture shifts with species-richness and climate across forests at continental scale. *Ecology Letters*, 25, 1442–1457.

Lilay, G.H., Thiebaut, N., du Mee, D., Assuncao, A.G.L., Schjoerring, J.K., Husted, S. et al. (2024) Linking the key physiological functions of essential micronutrients to their deficiency symptoms in plants. *New phytologist*, 242, 881–902.

Ma, S., He, F., Tian, D., Zou, D., Yan, Z., Yang, Y. et al. (2018) Variations and determinants of carbon content in plants: a global synthesis. *Biogeosciences*, 15, 693–702.

Maberly, S.C. & Gontero, B. (2018) Trade-offs and synergies in the structural and functional characteristics of leaves photosynthesizing in aquatic environments. In: *The Leaf: A Platform for Performing Photosynthesis* (eds. Adams III, WW & Terashima, I). Springer, pp. 307–343.

Makowski, D., Ben-Shachar, M. & Lüdecke, D. (2019) bayestestR: describing effects and their uncertainty, existence and significance within the bayesian framework. *Journal of Open Source Software*, 4, 1541.

Marschner, H. (2012) *Marschner's mineral nutrition of higher plants*.3rd ed. London, UK: Academic Press.

Moor, H., Rydin, H., Hylander, K., Nilsson, M.B., Lindborg, R. & Norberg, J. (2017) Towards a trait-based ecology of wetland vegetation. *Journal of Ecology*, 105, 1623–1635.

Pan, Y., Cieraad, E., Armstrong, J., Armstrong, W., Clarkson, B.R., Colmer, T.D. et al. (2020) Global patterns of the leaf economics spectrum in wetlands. *Nature Communications*, 11, 4519.

Author Manuscript

Pan, Y., García-Girón, J. & Iversen, L.L. (2023) Global change and plant-ecosystem functioning in freshwaters. *Trends in Plant Science*, 28, 646–660.

Pedersen, O., Colmer, T.D. & Sand-Jensen, K. (2013) Underwater photosynthesis of submerged plants - recent advances and methods. *Frontiers in Plant Science*, 4, 140.

Peñuelas, J., Fernández-Martínez, M., Ciais, P., Jou, D., Piao, S., Obersteiner, M. et al. (2019) The bioelements, the elementome, and the biogeochemical niche. *Ecology*, 100, e02652.

Piao, S., Ciais, P., Huang, Y., Shen, Z., Peng, S., Li, J. et al. (2010) The impacts of climate change on water resources and agriculture in China. *Nature*, 467, 43–51.

Rao, Q., Chen, J., Chou, Q., Ren, W., Cao, T., Zhang, M. et al. (2023) Linking trait network parameters with plant growth across light gradients and seasons. *Functional Ecology*, 37, 1732–1746.

Reich, P.B. (2014) The world-wide 'fast-slow' plant economics spectrum: a traits manifesto. *Journal of Ecology*, 102, 275–301.

Reich, P.B. & Oleksyn, J. (2004) Global patterns of plant leaf N and P in relation to temperature and latitude. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 11001–11006.

Reich, P.B., Oleksyn, J., Wright, I.J., Niklas, K.J., Hedin, L. & Elser, J.J. (2010) Evidence of a general 2/3-power law of scaling leaf nitrogen to phosphorus among major plant groups and biomes. *Proceedings of the Royal Society B: Biological Sciences*, 277, 877–883.

Reich, P.B., Walters, M.B. & Ellsworth, D.S. (1997) From tropics to tundra: global convergence in plant functioning. *Proceedings of the National Academy of Sciences of the United States of America*, 94, 13730–13734.

Sanchez, G. (2013) *PLS Path Modeling with R*. Trowchez Editions.

Sardans, J., Vallicrosa, H., Zuccarini, P., Farré-Armengol, G., Fernández-Martínez, M., Peguero, G. et al. (2021) Empirical support for the biogeochemical niche hypothesis in forest trees.

Author Manuscript

Nature Ecology & Evolution, 5, 184–194.

Schimel, D.S. (2003) All life is chemical. *Bioscience*, 53, 521–524.

Schoelynck, J. & Struyf, E. (2016) Silicon in aquatic vegetation. *Functional Ecology*, 30, 1323–1330.

Shen, M., Wang, S., Jiang, N., Sun, J., Cao, R., Ling, X. et al. (2022) Plant phenology changes and drivers on the Qinghai-Tibetan Plateau. *Nature Reviews Earth & Environment*, 3, 633–651.

Song, L., Wu, S., Ding, Y. & Luo, Y. (2021) *Change in Climate and Ecological Environment in China: 2021 Volume II Regional Impact, Vulnerability and Adaptation*. Science Press.

Sterner, R.W. & Elser, J.J. (2002) *Ecological stoichiometry: the biology of elements from molecules to the biosphere*. Princeton, NJ: Princenton University Press.

Tang, Z., Xu, W., Zhou, G., Bai, Y., Li, J., Tang, X. et al. (2018) Patterns of plant carbon, nitrogen, and phosphorus concentration in relation to productivity in China's terrestrial ecosystems. *Proceedings of the National Academy of Sciences of the United States of America*, 115, 4033–4038.

Wang, Z., Xia, C., Yu, D. & Wu, Z. (2015) Low-temperature induced leaf elements accumulation in aquatic macrophytes across Tibetan Plateau. *Ecological Engineering*, 75, 1–8.

Watanabe, T., Broadley, M.R., Jansen, S., White, P.J., Takada, J., Satake, K. et al. (2007) Evolutionary control of leaf element composition in plants. *New phytologist*, 174, 516-523.

Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F. et al. (2004) The worldwide leaf economics spectrum. *Nature*, 428, 821–827.

Xing, K., Niinemets, Ü., Rengel, Z., Onoda, Y., Xia, J., Chen, H.Y.H. et al. (2021) Global patterns of leaf construction traits and their covariation along climate and soil environmental gradients. *New phytologist*, 232, 1648–1660.

Yang, L., Zhao, H., Zuo, Z., Li, X., Yu, D. & Wang, Z. (2021) Generality and shifts in leaf trait relationships between alpine aquatic and terrestrial herbaceous plants on the Tibetan

Plateau. *Frontiers in Ecology and Evolution*, 9, 706237.

Yuan, M.M., Guo, X., Wu, L., Zhang, Y., Xiao, N., Ning, D. et al. (2021) Climate warming enhances microbial network complexity and stability. *Nature Climate Change*, 11, 343–348.

Zhao, N., Yu, G., He, N., Wang, Q., Guo, D., Zhang, X. et al. (2016) Coordinated pattern of multi-element variability in leaves and roots across Chinese forest biomes. *Global Ecology and Biogeography*, 25, 359–367.

Zuo, Z., Zhao, H., Yang, L., Lv, T., Li, X., Ma, F. et al. (2022) Salinity induces allometric accumulation of sulfur in plants and decouples plant nitrogen-sulfur correlation in alpine and arid wetlands. *Global Biogeochemical Cycles*, 36, e2022GB007372.

Author Manuscript

Author Manuscript

Figures legends

FIGURE 1 Conceptual frameworks for the shaping processes of plant elementome and the variations of plant element networks under stressful environments. (a), The realized plant elementomes are filtered by macroclimates and habitats from phylogeny-determined fundamental elementome. The bottom layer represents the species-specific physiological requirements for each element (E_1, E_2…) of plants, which determines the plant fundamental elementome. The middle layer indicates the influences of macroclimate (*e.g.*, temperature, precipitation and their combination). Macroclimate can determine the plant bioelemental concentrations by affecting the biochemical processes (*e.g.*, the Temperature-Plant Physiological Hypothesis, Reich & Oleksyn, 2004). The up layer is the realized elementome finally shaped by the habitat conditions (*e.g.*, nutrient supply and stresses). The nutrient supply determines the resource availability, while the environmental stresses limit the use of elements by plants. (b), The bivariate correlations among bioelements are spatialized into an element network (He et al., 2020). The variations in network topology towards stressful conditions can reflect the alterations in the element-element correlations, indicating the changes in bioelemental synergies to adapt to special habitats.

FIGURE 2 Stoichiometric characteristics of 17 bioelements of wetland plants. (a), Concentrations of 17 bioelements for overall observations (n = 1058) and different life-forms (submerged plants: n = 543; emergent plants: n = 475; floating-leaved plants: n = 40) of wetland plants. The concentrations of the left 11 bioelements refer to the left Y-axis, while the right 6 ones refer to the right Y-axis. Asterisks indicate statistical significance (P < 0.05) in leaf elements between different life-forms. (b), Coefficient of variation (CV, %) of 17 bioelements for overall observations and different life-forms of wetland plants (n = 17). (c), Linear regressions between

Author Manuscript

bioelemental concentration and its CV. (d), Linear regressions between bioelemental physiological requirement and its CV (n = 12).

FIGURE 3 Phylogenetic and environmental controls on 17 bioelements of wetland plants. (a), Variance partitioning for 17 bioelement variations using linear mixed effect model (n = 1028). Columns with different colors represent different variance components. (b), Linear regressions between bioelemental concentrations and phylogenetic variation (n = 17). Phylogenetic levels including the order level, order/family level, order/family/species level, family level and above, and species level and above. (c), Linear regressions between bioelemental concentrations and environmental variation (n = 17). Solid lines represent significant fits ($P < 0.05$).

FIGURE 4 Geographic and environmental drivers of bioelements of wetland plants (n = 1058)**.** (a)-(b), Heat map of the relationships between17 bioelements and environmental variables based on Bayesian phylogenetic linear mixed models. Colours indicate the probability of direction of the posterior distributions of the relationship between pairs of variables, where red and blue indicate positive and negative relationships respectively. Black dots indicate Bayes factors >1. Partial least squares path modeling (PLS-PM) for macroelements (c) and its standardized effects (d). PLS-PM for microelements (e) and its standardized effects (f). For panels (d) and (f), standardized effects are categorized into standardized direct effects and standardized indirect effects. Only significant pathways ($P < 0.05$) are showed in these models. The values adjacent to arrows are standardized path coefficients and the arrow width is proportional to the intensity of relationship. R^2 indicates the explained variance of dependent variable by the model.

FIGURE 5 Shifts in leaf element networks (LENs) and network parameters along environmental gradients of wetland plants. (a), The network topologies of LENs for all data (n = 1058) and different temperature and salinity gradients. High-GST (n = 346), middle-GST (n = 498) and low-GST (n = 214) represent 20-30 ℃, 10-20 ℃ and 0-10 ℃ of growing season mean temperature (GST), respectively. Fresh water (n = 757), brackish water (n = 202), moderate saline

Author Manuscript

water (n = 69) and saline water (n = 30) represent 0-0.5‰, 0.5-2‰, 2-5‰ and >5‰ of water salinity, respectively. Different colors in each network represent different modules. The size of the node represents its degree, and the width of the edge represents the strength of positive (red lines) and negative (blue lines) correlations. (b), The variations in network overall parameters (modularity, edge density, average clustering coefficient, average path length, diameter and vulnerability) across temperature and salinity gradients, respectively.

FIGURE 6 Highlights for explaining the shaping mechanisms of wetland plant elementome. (a), With increasing bioelemental concentrations from microelements to macroelements, the elements show more stable (red line) and less variable (brown line), and the driving factors for the variations of bioelements shift from environmental modification (black line) to evolutionary constraint (blue line). (b), The underlying environmental controls of macroelements and microelements are divergent. Nutrient supplies in sediment and water are the primary controlling factors in determining the patterns of microelements, but climate play more important roles in the variations in macroelements than microelements. (c), The topologies of leaf element network vary along environmental gradients. More intense stresses decrease network connectivity, complexity and stability, which means that the synergies among bioelements become weaker in stressful

habitats. Elements with high importance consistently maintain their paramount importance in plant elementome across environmental gradients.

GA text:

Our findings reveal that leaf bioelements with higher concentration were more stable and evolutionary constrained. We proposed a Stability of Well-coordinated Elements Hypothesis, suggesting that bioelements that coordinate well in driving certain physiological functions constrain each other, thus maintaining relatively stable ratios in plants. Our research filled the gap in study of wetland plant elementome beyond C, N and P, and provided new evidences to predict plant adaptations and plant-environment interactions in regions sensitive to climate change.

Author Manuscript

Author Manuscript




Figure 1.tif

Author Manuscript




Figure 2.tif

Author Manuscript




Figure 3.tif

Author Manuscript




Figure 4.tif

Author Manuscript




Figure 5.tif

Author Manuscript




Figure 6.tif